## All required fields (marked with \*) and signatures must be completed before submitting this form by fax or email.

Fax referral to: 1–888–668–2143 | Email referral to: AmplifyAssist@orsinihc.com Call AmplifyAssist at 888–668–4198 for more information | Hours: 8 am–6 pm CT, Monday–Friday

## **AmplifyAssist™ Enrollment Form**





		1	. PATIENT INFO	RMATION				
First Name:		Middle Initial:	*Last Na	mo:		Date of Birth://		
		Other: (P				/		
			•	enification purposes.)				
	-	Spanish Other:						
•						•		
						Communication: Phone Text Ema		
				•		State: Zip Code:		
Representative/Caregiver Name: Relationship to Patient:								
Primary Phone	Number:			Mobile Best Time t	o Call:	AM PM No preference		
		2.	. CLINICAL INFO	DRMATION				
Primary Diagno	osis Code: 🗌 Ple	ase see attached clinical information	for the information r	equested below.				
	☐ E75	5.24 = Niemann-Pick disease	E75.242	= Niemann-Pick dise	ase type C			
	☐ E75	5.249 = Niemann-Pick disease, unspe	ecified 🗌 Other: _					
Is/Has the patient been treated with miglustat?   Yes   No Dose:   Parti						icipant/Participated in Zevra Expanded Access Progra		
•		•			·	· · · · · · · · · · · · · · · · · · ·		
	3. 11	ISURANCE INFORMATION (F	lease provide t	ront and back c	ppy of in	surance card(s))		
rimary Insuran	ce Commer	cial Medicare Medicaid	Secondary Insurance					
surance Carri	er Name:		Insurance Carrier Name:					
Phone Number:				Phone Number:				
Employer Grp/Issuer if available: Empl					Employer Grp/Issuer if available:			
hone Number:			Phone Number:					
)#:		Group#:	ID#: Group#:					
		·						
		Group#:						
		PCN#:		•				
		4 055105	AND DDEACDID	FION INFORMAT	ION			
				TION INFORMAT				
						*NPI:*		
	•		*City:					
				#:				
Phone:				.,.				
	Completing This	Form:		*Email	:			
Office Contact Recommended	Dosing Guidelir	nes: Please see <u>Prescribing</u>	*Patient Weigl	*Email nt kg or		*Dispense:		
Office Contact	Dosing Guidelir		*Patient Weigl *Prescription:	ntkg or		•		
Office Contact Recommended Information fo	l <b>Dosing Guidelir</b> r dosing consider	nes: Please see <u>Prescribing</u>	*Prescription:  47 mg (NE	nt kg or OC: 72542-147-01)		•		
Office Contact Recommended Information fo	l <b>Dosing Guidelir</b> r dosing consider	nes: Please see <u>Prescribing</u> rations in special populations.	*Prescription:  47 mg (NE  62 mg (NE	nt kg or OC: 72542-147-01) OC: 72542-162-01)		#90 capsules Other:*Refills:		
Office Contact Recommended Information for Patient Body W	Dosing Guidelir r dosing consider	res: Please see <u>Prescribing</u> rations in special populations.  Recommended Dosage	*Prescription:  47 mg (NE) 62 mg (NE) 93 mg (NE)	kg or DC: 72542-147-01) DC: 72542-162-01) DC: 72542-193-01)		#90 capsules Other:*Refills:		
Office Contact Recommended Information for Patient Body W 8-15 kg	Dosing Guidelir r dosing consider /eight 17.6-33 lb	Recommended Dosage 47 mg three times a day	*Prescription:	kg or		#90 capsules Other:  *Refills: Other:  #90 capsules Other:  *SIG:		
Office Contact Recommended Information fo Patient Body W 8-15 kg >15-30 kg	Dosing Guidelir r dosing consider /eight 17.6-33 lb >33-66 lb	Recommended Dosage 47 mg three times a day 62 mg three times a day	*Prescription:	kg or DC: 72542-147-01) DC: 72542-162-01) DC: 72542-193-01)		#90 capsules Other:  *Refills:  #90 capsules Other:  *SIG:  Take one capsule by mouth three times d		
Product capsules	Posing Guideling dosing consider dosing consider designs and designs are designs and designs and designs and designs and designs are designs and designs and designs are designs and designs and designs are designs and designs and designs and designs are designs are designs are designs and designs are designs are designs are designs and designs are designs are designs are designs are designs are designs are desig	rations in special populations.  Recommended Dosage  47 mg three times a day  62 mg three times a day  93 mg three times a day  124 mg three times a day  whole or the contents of the capsule a soft food, or added to water to	*Prescription:	kg or		#90 capsules Other:  *Refills: Other:		
Recommended information for a second with the	J Dosing Guideling of dosing consider dosing consider designs of the dosing consider designs of the dosing consider dosing the dosin	rations in special populations.  Recommended Dosage  47 mg three times a day  62 mg three times a day  93 mg three times a day  124 mg three times a day  whole or the contents of the capsule a soft food, or added to water to	*Prescription:	kg or	lb	#90 capsules Other: *Refills:		